

**ITEM 1**

**ReSoMal, 42g sachet for 1 litre/CAR-100**

General Description:

ReSoMal (ORS for severely malnourished children), 42g sachet to be diluted in 1 litre of purified/boiled and cooled water, carton of 100 sachets.

Technical Specifications:

Contents per sachet of 42 grams:

Sucrose: 25g

Glucose Anhydrous: 10g

Sodium Chloride: 1.75g

Trisodium Citrate, dihydrate: 1.45g

Potassium Chloride: 2.54g

Tripotassium Citrate: 0.65g

Magnesium Chloride Anhydrous: 0.61g

Zinc Acetate: 0.0656g

Copper Sulphate Anhydrous: 0.0112g

Osmolarity: 300 mmol/l

All the ingredients must comply with one of the pharmacopeias:

BP, Ph.Eur, Ph.Int, USP.

Instructions for use

One sachet of 42.1 grams is to be diluted in 1 liter of purified/boiled and cooled water which will give 1 liter of liquid ReSoMal.

The reconstituted solution has to be consumed immediately or used within 24 hours if stored in a refrigerator.

Target population (beneficiaries: type and number of persons concerned):

Children 6-59m old suffering from Dehydration AND Severe Acute Malnutrition

Should not be used for cholera patients.

## ITEM 2

### ReSoMal, 42g sachet for 1 litre/CAR-50

#### General Description:

ReSoMal (ORS for severely malnourished children), 42g sachet to be diluted in 1 litre of purified/boiled and cooled water, carton of 50 sachets.

#### Technical Specifications:

Contents per sachet of 42 grams:

Sucrose: 25g

Glucose Anhydrous: 10g

Sodium Chloride: 1.75g

Trisodium Citrate, dihydrate: 1.45g

Potassium Chloride: 2.54g

Tripotassium Citrate: 0.65g

Magnesium Chloride Anhydrous: 0.61g

Zinc Acetate: 0.0656g

Copper Sulphate Anhydrous: 0.0112g

Osmolarity: 300 mmol/l

All the ingredients must comply with one of the pharmacopeias:

BP, Ph.Eur, Ph.Int, USP.

#### Instructions for use

One sachet of 42.1 grams is to be diluted in 1 liter of purified/boiled and cooled water which will give 1 liter of liquid ReSoMal.

The reconstituted solution has to be consumed immediately or used within 24 hours if stored in a refrigerator.

Target population (beneficiaries: type and number of persons concerned):

Children 6-59m old suffering from Dehydration AND Severe Acute Malnutrition

Should not be used for cholera patients.